

Patient

Name: _____ Phone #: _____ Patient ID #: _____

Fasting Status: **12 HOURS** File ID: _____ Gender: **MALE** Birthdate: _____ Age: _____

Height: **5 ft 10 in** Weight: **210 lbs** BMI: **30.1** Prev. BMI: **31.6**

Specimen

Collection Time: **8:50 am** Specimen ID: **17032900596**

Collection Date: **3/28/2017** Report Type: **COMPLETE**

Received Date: **3/29/2017** Report Date: **3/30/2017**

Provider

Requesting Provider: **DANIEL MELVILLE, MD**
DESTINATION HEALTH
900 E. SOUTHLAKE BLVD, SUITE 200
SOUTHLAKE, TX 76092

Client ID: _____

Laboratory Test	Notes	High Risk	Intermediate Risk	Optimal	High Risk Range	Intermediate Risk Range	Optimal Range	Previous Results 11/21/2016
-----------------	-------	-----------	-------------------	---------	-----------------	-------------------------	---------------	--------------------------------

Metabolic	HbA1c (%)			5.6	≥ 6.5	5.7 - 6.4	≤ 5.6	5.6
	Estimated Average Glucose (mg/dL) (calculated)			114.0	≥ 139.9	116.9 - 139.8	≤ 116.8	114.0
	25-hydroxy-Vitamin D (ng/mL)			46	< 20	20 - 29	30 - 100	34
	Uric Acid (mg/dL)			6.3	≥ 8.0	7.0 - 7.9	2.0 - 6.9	
	TSH (μIU/mL)			2.41	< 0.27 or > 4.20		0.27 - 4.20	
	Homocysteine (μmol/L)	14			> 13	11 - 13	< 11	12
	Vitamin B ₁₂ (pg/mL)			615	< 211	211 - 400	> 400	
	RBC Folate (ng/mL)			> 1396	< 700	700 - 750	> 750	
	CoQ10 (μg/mL) ¹			1.82	< 1.11	1.11 - 2.00	> 2.00 Target of therapy for patients on statins is > 2.0 μg/mL	

TSH is analyzed using reagents from Roche Diagnostics by electrochemiluminescence immunoassay. These values should not be used in conjunction with values from other reagent manufacturers or methodologies.

Renal	Cystatin C (mg/L)			0.85	≥ 1.04	0.96 - 1.03	≤ 0.95	
	Estimated Glomerular Filtration Rate (eGFR, mL/min/1.73m ²)			110	< 60	60 - 89	> 89	
	Microalbumin (urine) (mg albumin/g of creatinine)			7	≥ 30		≤ 29	
	Creatinine, serum (mg/dL)			0.9	> 1.2		0.7 - 1.2	

Lab Notes:

